

REPORT DOCUMENTATION PAGE				<i>Form Approved</i> <i>OMB No. 0704-0188</i>	
<small>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</small> PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01-NOV-2007		2. REPORT TYPE Journal Article POSTPRINT		3. DATES COVERED (From - To) xx-XXX-200x -- 30-SEP-2005	
4. TITLE AND SUBTITLE Conversion of 2-(4-carboxyphenyl)-6-nitrobenzothiazole to 4-(6-amino-5-hydroxybenzothiazol-2-yl)benzoic acid by a recombinant E. coli strain				5a. CONTRACT NUMBER FA4819-07-D-0001	
				5b. GRANT NUMBER 	
				5c. PROGRAM ELEMENT NUMBER 62102F	
6. AUTHOR(S) Nadeau, Lloyd J.; Spain, Jim C.; Kannan, Ramamurthi; Tan, Loon-Seng				5d. PROJECT NUMBER 4915	
				5e. TASK NUMBER E2	
				5f. WORK UNIT NUMBER 4915E30A	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory Materials and Manufacturing Directorate 139 Barnes Drive, Suite 2 Tyndall Air Force Base, FL 32403-5323				8. PERFORMING ORGANIZATION REPORT NUMBER 	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory Materials and Manufacturing Directorate 139 Barnes Drive, Suite 2 Tyndall Air Force Base, FL 32403-5323				10. SPONSOR/MONITOR'S ACRONYM(S) AFRL/RXQL	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-ML-TY-TP-2005-4605	
12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Statement A: Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES Published in Chem Comm Communication (2006), 564-565. Pub. The Royal Society of Chemistry.					
14. ABSTRACT E. coli C43(DE3)pNbzAHabA, expressing nitoreductase and mutase enzymes, converts 2-(4-carboxyphenyl)-6-nitrobenzothiazole to 4-(6-amino-5-hydroxybenzothiazol-2-yl)benzoic acid.					
15. SUBJECT TERMS 					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON Glenn Johnson
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (Include area code)

Reset

Conversion of 2-(4-carboxyphenyl)-6-nitrobenzothiazole to 4-(6-amino-5-hydroxybenzothiazol-2-yl)benzoic acid by a recombinant *E. coli* strain†

Lloyd J. Nadeau,^a Jim C. Spain,^{*b} Ramamurthi Kannan^c and Loon-Seng Tan^c

Received (in Cambridge, MA, USA) 10th November 2005, Accepted 28th November 2005

First published as an Advance Article on the web 21st December 2005

DOI: 10.1039/b516032d

E. coli C43(DE3)pNbzAHabA, expressing nitroreductase and mutase enzymes, converts 2-(4-carboxyphenyl)-6-nitrobenzothiazole to 4-(6-amino-5-hydroxybenzothiazol-2-yl)benzoic acid.


4,6-Diaminoresorcinol (**1**) and 2,5-diamino-1,4-benzenedithiol (**2**) (usually stored as dihydrochloride salts) are key co-monomers for the synthesis of strong, thermally resistant rigid-rod poly(*p*-phenylenebenzobisoxazole) (PBO) and poly(*p*-phenylenebenzobisthiazole) (PBT) polymers (Scheme 1) for lightweight structural, nonlinear optical and electronic applications.^{1,2} More recently, PBO has been considered for use in the proton-exchange membranes of fuel cells.³ While an AB-monomer 4-[5-amino-6-hydroxybenzoxazol-2-yl]benzoic acid (**3**) has been prepared and utilized in the synthesis of the corresponding PBO polymer,⁴ the analogous AB-monomer for PBT (**4**) has not been synthesized. In addition to having an intrinsically perfect stoichiometry that helps to promote high molecular weight polymers in polycondensation processes, AB-monomers are also useful starting materials for the synthesis of AB diblock copolymers, ABA triblock copolymers and star polymers. They can also be grafted onto appropriately functionalized surfaces. The availability of the above aminophenol derivative is limited because the synthesis is complex and yields are low. Recent advances in biologically converting aromatic nitro compounds to the corresponding *o*-aminophenols suggest the possibility of synthesizing novel AB-monomers such as 4-(6-amino-5-hydroxybenzothiazol-2-yl)benzoic acid (**5**) and in turn, the corresponding rigid-rod polymer that is a hybrid of both PBO and PBT with respect to the chemical structure.

Although the requisite starting nitro compounds (either as carboxylic acid, **7a** or ethyl ester **7b**) are relatively simple molecules, they have not been reported to our knowledge. In principle, **5** could be synthesized in two steps: (i) a condensation reaction between 2-aminothiophenol and 4-carboxybenzaldehyde, followed by (ii) nitration of the resulting 2-(4-carboxyphenyl)benzothiazole (**7a**). In practice, the poor solubility of **7a** and **8a** has

necessitated additional esterification/de-esterification steps in order to rigorously establish the identity and purity of the target nitrobenzothiazolecarboxylic acid. The detailed preparative procedures for **7a–8b** are provided as supplementary information.

Previously, we reported that nitroarenes are converted to *o*-aminophenols by nitroreductase and mutase enzymes.⁵ The nitroreductase reduces nitroaromatic compounds to hydroxylaminoarenes⁶ and hydroxylaminobenzene mutase catalyzes a regio-specific reaction converting hydroxylaminoarenes to the corresponding *o*-aminophenols by an intramolecular transfer of the hydroxyl group.⁷ Furthermore, an *E. coli* containing nitroreductase and mutase converts simple nitroarenes regio-specifically to *o*-aminophenols at high yields.⁸

In preliminary experiments, cells of *E. coli* C43(DE3)-pNbzAHabA (*i.e.* strain JS995) were grown and induced as described previously.⁸ 2-(4-Carboxyphenyl)-6-nitrobenzothiazole **7a** (49 μ M) was incubated with the cells and HPLC analysis of the reaction mixture revealed a transformation rate for the parent compound of 0.9 nmoles min⁻¹ mg⁻¹ protein and the accumulation of one product (Fig. 1). LC/MS (+APCI) analysis of the product revealed a 287 *m/z* as expected of an aminophenol. The conversion efficiency was 100%. The transformation was scaled up



Scheme 1 Chemical structures of pertinent monomers, polymers, substrate and intermediates.

^aAFRL/MLQL, Tyndall AFB, 139 Barnes Dr., Building 1117, FL 32403, USA. E-mail: lloyd.nadeau@tyndall.af.mil; Fax: 850-283-6090; Tel: 850-283-6018

^bGeorgia Institute of Technology, 311 Ferst Dr., Atlanta, GA 30332-0512, USA. E-mail: jspain@ce.gatech.edu; Fax: 404-894-8266; Tel: 404-894-0628

^cAFRL/MLBP, Wright-Patterson AFB, 2941 Hobson Way, B654 R309, OH 45433-7750, USA. E-mail: loon-seng.tan@wpafb.af.mil; Fax: 937-255-9157; Tel: 937-255-9141

† Electronic supplementary information (ESI) available: experimental section. See DOI: 10.1039/b516032d




Fig. 1 Transformation of 2-(4-carboxyphenyl)-6-nitrobenzothiazole to 4-(6-amino-5-hydroxybenzothiazol-2-yl)benzoic acid by *E. coli* C43(DE3)pNbzAHabA.

to a 2 L bioreactor containing M9 medium⁹ (30 °C) supplemented with sorbitol (20 g/L), ampicillin (100 μg/L) and isopropylthio-β-D-galactoside (1 mM). Induced cells were added to the bioreactor ($A_{600} = 8.0$) and **7a** (5 mM) dissolved in NH_4OH (2 N) at 65 °C was delivered repeatedly to the reactor. The disappearance of reactant and the accumulation of product were monitored by HPLC. Over a 2 hour period, 200 mg of **7a** were converted. The cells were removed by centrifugation and the product was precipitated by adjusting the pH of the supernatant to 2.7 with HCl. Green crystals were recovered by centrifugation. The crystals were dissolved in NH_4OH , filtered and recrystallized by lowering the pH to 2.7. The pelleted crystals were dried overnight under vacuum, washed with water, and then with acetone. The melting point was 326–328 °C.

The proton NMR spectrum (270 MHz) of the purified compound in DMSO-d_6 showed that in the aromatic proton region, there were three singlet peaks at δ (ppm) 7.117, 7.309, and 8.012 at relative intensities of 1 : 1 : 4. The fact that only two distinct singlets were observed from the two protons on the phenyl ring with tetrasubstitution ruled out the isomeric structure **6** or mixture of **5** and **6** (Fig. 2). The FT-IR (KBr) spectrum (ESI) is consistent with the NMR data, indicating the presence of $\nu(\text{C}=\text{O})$ of carboxylic acid at 1692 cm^{-1} and a strong, broad band centered at $\sim 3431\text{ cm}^{-1}$ that is attributable to the hydroxyl-group vibrations of the carboxylic acid and the phenol moieties. The symmetrical and asymmetrical NH_2 stretches, typically detected as a doublet at ~ 3400 and $\sim 3500\text{ cm}^{-1}$ respectively, are most likely hidden beneath the broad $\nu(\text{OH})$ band. Electron-impact mass spectroscopy gave a molecular ion with $m/z = 285.96$ (100% relative abundance). Thus, all the available spectroscopic data confirm the structure of the product as 4-(6-amino-5-hydroxybenzothiazol-2-yl)benzoic acid, **5**.

The biocatalyst, *E. coli* C43(DE3)pNbzAHabA converts 2-(4-carboxyphenyl)-6-nitrobenzothiazole to a potentially useful *ortho*-aminophenolic synthon for the synthesis of novel polymers. Our previous work indicated that the combination of the reductase and mutase enzymes could catalyze the transformation of very simple nitroaromatic compounds to the corresponding *ortho*-aminophenols. The results presented here indicate that the biocatalyst can transform more complex and potentially useful nitroaromatic compounds stoichiometrically to the *ortho*-aminophenols. Such




Fig. 2 Proton NMR spectrum of aminophenol product in DMSO-d_6 . For clarity, only the aromatic proton region is shown.

conversions using traditional organic chemistry would be prohibitively complex and expensive. We are currently exploring the ability of the biocatalyst to transform a variety of other complex molecules.

We thank Michael Matuszewski and Ann Dombroskie for their assistance in the preparation of the nitrobenzothiazole-based substrates, Glenn Johnson and Ion Ghiriviga for helpful discussions, and the Air Force Office of Scientific Research for funding.

Notes and references

- X.-D. Hu, S. E. Jenkins, B. G. Min, M. B. Polk and S. Kumar, *Macromol. Mater. Eng.*, 2003, **288**, 823–843; R. C. Evers, F. E. Arnold and T. E. Helminiak, *Macromolecules*, 1981, **14**, 925–930; S. Kumar, T. D. Dang, F. E. Arnold, A. R. Bhattacharyya, B. G. Min, X. Zhang, R. A. Vaia, C. Park, W. W. Adams, R. H. Hauge, R. E. Smalley, S. Ramesh and P. A. Willis, *Macromolecules*, 2003, **35**, 9039–9043; Y.-H. So, *Acc. Chem. Res.*, 2001, **34**, 753–758.
- Y.-H. So, *Prog. Polym. Sci.*, 2000, **25**, 137–157; S. A. Jenekhe, L. R. de Paor, X. L. Chen and R. M. Tarkka, *Chem. Mater.*, 1996, **8**, 2401–2404; S. A. Jenekhe, J. A. Osaheni, J. S. Meth and H. Vanherzeele, *Chem. Mater.*, 1992, **4**, 683–687.
- S.-Y. Park, H. Koerner, S. Putthanarat, R. Ozisik, S. Juhl, B. L. Farmer and R. K. Eby, *Polymer*, 2004, **45**, 49–59.
- J. P. Chen and W. A. Feld, *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)*, 1990, **31**, 681–682; M. Dotrong, M. H. Dotrong and R. C. Evers, *J. Polym. Sci., Part A: Polym. Chem.*, 1997, **35**, 3451–3456.
- S. F. Nishino and J. C. Spain, *Appl. Environ. Microbiol.*, 1993, **59**, 2520–2525; A. Schenzle, H. Lenke, P. Fischer, P. A. Williams and H.-J. Knackmuss, *Appl. Environ. Microbiol.*, 1997, **63**, 1421–1427; H.-S. Park and H.-S. Kim, *J. Bacteriol.*, 2000, **182**, 573–580; Y. Hasegawa, T. Muraki, T. Tokuyama, H. Iwaki, M. Tatsuno and P. C. K. Lau, *FEMS Microbiol. Lett.*, 2000, **190**, 185–190; T. Spiess, F. Desiere, P. Fischer, J. C. Spain, H.-J. Knackmuss and H. Lenke, *Appl. Environ. Microbiol.*, 1998, **64**, 446–452.
- H. Nivinskas, R. L. Koder, Z. Anusevicius, J. Sarlauskas, A.-F. Miller and N. Cenas, *Arch. Biochem. Biophys.*, 2001, **385**, 170–178; C. C. Somerville, S. F. Nishino and J. C. Spain, *J. Bacteriol.*, 1995, **177**, 3837–3842.
- L. J. Nadeau, Z. He and J. C. Spain, *Appl. Environ. Microbiol.*, 2003, **69**, 2786–2793.
- V. Kadiyala, L. J. Nadeau and J. C. Spain, *Appl. Environ. Microbiol.*, 2003, **69**, 6520–6526; L. J. Nadeau, Z. He and J. C. Spain, *J. Ind. Microbiol. Biotechnol.*, 2000, **24**, 301–305.
- J. Sambrook, E. F. Fritsch and T. Maniatis, *Molecular cloning: a laboratory manual*, 2nd edn., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.